

**Brain injury occurring during or soon after birth: annual incidence and rates of brain injuries to monitor progress against the national maternity ambition
2018 and 2019 national data**

Authors: Dr Chris Gale, Miss Kayleigh Ougham, Ms Sena Jawad, Dr Sabita Uthaya, Professor Neena Modi

Affiliation: Neonatal Data Analysis Unit, Imperial College London, Chelsea and Westminster Hospital campus, SW10 9NH

Date of production: 28th January 2021

Contents

	Page
1 Context	3
2 Methods	4
2.1 Defining brain injuries occurring during or soon after birth	4
2.2 Numerator data source: National Neonatal Research Database	5
2.3 Denominator Data Sources: National live births	6
2.4 Statistical Methods	7
2.5 Erratum relating to data 2012-2017	8
3 Results	9
3.1 National rates of brain injuries	9
3.1.1 Rates of brain injuries in England: all gestational ages	9
3.1.2 Rates of brain injuries in England: term infants	12
3.1.3 Rates of brain injuries in England: preterm infants	15
3.1.4 Rates of brain injuries in England by medical condition	18
Appendix 1: Defining brain injuries	24

1 Context

In November 2015, the Secretary of State for Health announced a national ambition to halve the annual rates of stillbirth, neonatal death, maternal death and brain injuries occurring during or soon after birth in England.

The Department of Health recognised that there was a lack of consensus in relation to the definition of *brain injury occurring during or soon after birth*, and limited data sources for measuring such brain injuries. Therefore the Department of Health commissioned a piece of work to define *brain injury occurring during or soon after birth*, and using the National Neonatal Research Database (NNRD) established a baseline for 2010 and calculated annual rates up to 2017. These data have been previously published^{1 2 3}.

The Department of Health and Social Care want to build on this previous work, specifically to track progress towards the National Maternity Ambition to halve the rate of *brain injuries occurring during or soon after birth* by 2025, with a 20% reduction by 2020⁴.

In this follow-up report we provide the following rates for 2018 and 2019 in England:

- (a) Annual rates of *brain injuries occurring during or soon after birth* for all gestational ages, and for term and preterm infants separately.**
- (b) Annual rates of individual conditions that contribute to *brain injuries occurring during or soon after birth*.**

¹ Brain injury occurring during or soon after birth: a report for the national maternity ambition commissioned by the Department of Health

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/662974/Report_on_brain_injury_occurring_during_or_soon_after_birth.pdf

² Gale C, Statnikov Y, Jawad S On behalf of the Brain Injuries expert working group, et al., Neonatal brain injuries in England: population-based incidence derived from routinely recorded clinical data held in the National Neonatal Research Database. *Archives of Disease in Childhood - Fetal and Neonatal Edition*

<https://fn.bmj.com/content/early/2017/10/22/archdischild-2017-313707>

³ Gale C, Jeyakumaran D, Ougham K, Jawad S, Uthaya S, Modi N. Brain injury occurring during or soon after birth: annual incidence and rates of brain injuries to monitor progress against the national maternity ambition. 2016 and 2017 data

<https://www.imperial.ac.uk/media/imperial-college/medicine/dept-medicine/infectious-diseases/neonatology/Brain-injury-occurring-during-or-soon-after-birth-120419-V3.3.pdf>

⁴ Safer maternity care: progress and next steps. Department of Health, November 2017

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/662969/Safer_maternity_care_-_progress_and_next_steps.pdf

2 Methods

2.1 Defining brain injuries occurring during or soon after birth

The definition for *brain injuries occurring during or soon after birth* for the national maternity ambition was agreed by an expert group as previously described⁵. This definition is as follows:

- **Population: *all babies admitted to a neonatal unit in England***
- **Time period after birth: *all brain injuries that are detected during the neonatal unit stay***
- **Conditions to be included:**
 - a. *Infants with a diagnosis consistent with **hypoxic ischaemic encephalopathy**: term and near-term infants only*
 - b. *Infants with a diagnosis of **intracranial haemorrhage, perinatal stroke, hypoxic ischaemic encephalopathy (HIE), central nervous system infection, and kernicterus (bilirubin encephalopathy)**: all infants*
 - c. ***preterm white matter disease (periventricular leukomalacia)**: preterm infants only*
 - d. *Infants with a recorded **seizure**: all infants*
- **Denominator: *all live births in England* to be used as the denominator for calculating the annual rate of “brain injuries occurring during or soon after birth”**
- **Exclusions:** a consensus decision⁴ was made to present data before and after exclusion of infants with the following conditions leading to brain injury prior to birth: ***congenital encephalopathies (including inborn errors of metabolism), congenital infections and congenital brain abnormalities***

Further descriptive information and the NNRD data fields used to determine *brain injuries occurring during or soon after birth* and exclusions are shown in Appendix 1.

⁵ Brain injury occurring during or soon after birth: a report for the national maternity ambition commissioned by the Department of Health
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/662974/Report_on_brain_injury_occurring_during_or_soon_after_birth.pdf

2.2 Numerator data source: National Neonatal Research Database

The Department of Health and Social Care commissioned the National Neonatal Research Database (NNRD) as the numerator source for the calculation of rates of *brain injuries occurring during or soon after birth* for the national maternity ambition.

The NNRD is formed from summary electronic patient data entered on all admissions to National Health Service (NHS) neonatal units in England. Data in the NNRD have undergone processing to identify duplicates, out of range values, internal inconsistencies, and other potentially erroneous entries. In addition to these internal processes, feedback and quality assurance checks are undertaken with clinicians for key items. Data in the NNRD are merged across multiple patient episodes, if necessary for the purposes of an analysis.

All neonatal units (currently 100% of NHS neonatal units in England), that contribute data to the NNRD form the UK Neonatal Collaborative, UKNC. Approximately 450 predefined data items, the Neonatal Data Set⁶, are extracted quarterly from these electronic patient records to form the NNRD. The Neonatal Data Set is an approved NHS Information Standard (ISB1595) and the NNRD is a national Information Asset.

National counts of *brain injuries occurring during or soon after birth* are calculated using a cohort of infants who receive at least one episode of care in a NHS neonatal unit in England. As part of the NNRD data cleaning process, babies with missing gestational age or missing birth year are excluded from the analysis data set. To extract counts for term and preterm babies, gestational ages of ≥ 37 weeks and < 37 weeks respectively were applied.

⁶http://www.datadictionary.nhs.uk/data_dictionary/messages/clinical_data_sets/data_sets/national_neonatal_data_set/national_neonatal_data_set_-_episodic_and_daily_care_fr.asp?shownav=0

2.3 Denominator data sources: National live births

To calculate national rates for 2018 and 2019, denominator data from the Office for National Statistics (ONS) Birth Summary Tables for England⁷, describing total live births by gestational age (in weeks) for 2018 and 2019 where mothers' usual residence is England, were used. These data were also used as the denominator when calculating rates of medical conditions leading to *brain injuries occurring during or soon after birth* where gestational age is specified. Births were extracted from a dataset containing birth registrations linked to their corresponding notification and infants missing gestational age and where birth weight was inconsistent with gestational age were excluded. As a result, lower numbers of live births are reported for these analyses than for analyses that include all infants.

⁷ Office for National Statistics, last accessed 14th December 2020, <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/12561livebirthsstillbirthsandneonataldeathsbygestationalageinengland2010to2018neonataldeathsand2010to2019livebirthsandstillbirths>

2.4 Statistical methods

We present rates of brain injuries per 1,000 live births unless otherwise stated. For years 2012-2017, rates of brain injuries were calculated as described previously^{8 9}.

Confidence intervals (CI) for rates have been calculated using the following method:

- 1) Calculate error factor (EF):

$$EF = e^{(1.96/\sqrt{D})}$$

where D = number of brain injuries

- 2) Calculate lower and upper CI:

$$\text{Lower CI} = \text{Rate} \div EF$$

$$\text{Upper CI} = \text{Rate} \times EF$$

⁸ Gale C, Statnikov Y, Jawad S et al on behalf of the Brain Injuries expert working group, Neonatal brain injuries in England: population-based incidence derived from routinely recorded clinical data held in the National Neonatal Research Database. *Archives of Disease in Childhood - Fetal and Neonatal Edition* <https://fn.bmj.com/content/early/2017/10/22/archdischild-2017-313707>

⁹ Gale C, Jeyakumaran D, Ougham K, Jawad S, Uthaya S, Modi N. Brain injury occurring during or soon after birth: annual incidence and rates of brain injuries to monitor progress against the national maternity ambition. 2016 and 2017 data. Last accessed 18th January 2021 <https://www.imperial.ac.uk/media/imperial-college/medicine/dept-medicine/infectious-diseases/neonatology/Brain-injury-occurring-during-or-soon-after-birth-120419-V3.3.pdf>

2.5 Erratum relating to data 2012-2017

An error in data extraction meant that rates of brain injuries previously reported for 2010-2017 require the following corrections:

Erroneous data extraction resulted in:

1. Incorrect inclusion in the measure *brain injuries occurring during or soon after birth* of babies with Grade 1 Hypoxic Ischaemic Encephalopathy (HIE) who did not receive therapeutic hypothermia
2. Incorrect exclusion of babies with central nervous system infection recorded after 48 hours after birth from the measure *brain injuries occurring during or soon after birth*

These errors have been corrected and rates of *brain injuries occurring during or soon after birth* for 2012-2017 have been recalculated using corrected data in the table and figures in this report.

3 Results

3.1 National rates of brain injuries

3.1.1 Rates of brain injuries in England: all gestational ages

Numerator:	Annual number of infants who received at least one episode of care within a neonatal unit in England with a <i>brain injury occurring during or soon after birth</i> , with and without exclusions, obtained from the NNRD.
Denominator:	Annual number of live births in England obtained from ONS Birth Summary Tables ¹⁰ .

Table 1 presents the incidence and crude rates per 1,000 live births of brain injuries, with and without exclusions of infants with conditions leading to brain injury prior to birth, for years 2012 to 2019. Figure 1 displays the crude rates of brain injuries per 1,000 live births prior to exclusions.


¹⁰ Office for National Statistics, last accessed 14th December 2020, <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/12561livebirthsstillbirthsandneonataldeathsbygestationalageinengland2010to2018neonataldeathsand2010to2019livebirthsandstillbirths>

Table 1: Infants in England (all gestational ages) with a diagnosis of brain injury, before and after exclusion of infants with conditions leading to brain injury prior to birth.

Year	Infants recorded in the NNRD	Infants recorded in the NNRD with GA available (%)	Live births in England	Before exclusion of infants with conditions leading to brain injury prior to birth		After exclusion of infants with conditions leading to brain injury prior to birth		
				Infants with brain injury	Rate of brain injuries per 1000 live births (95%CI)	Exclusions	Infants with brain injury	Rate of brain injuries per 1000 live births (95% CI)
2012	78,980	78,952 (99.96)	694,241	2,950	4.25 (4.10,4.41)	76	2,874	4.14 (3.99, 4.29)
2013	80,222	80,199 (99.97)	664,517	2,966	4.46 (4.31,4.63)	62	2,904	4.37 (4.21, 4.53)
2014	85,013	84,981 (99.96)	661,496	3,097	4.68 (4.52,4.85)	56	3,041	4.60 (4.44,4.76)
2015	88,931	88,785 (99.84)	664,399	3,055	4.60 (4.44,4.76)	54	3,001	4.52 (4.36, 4.68)
2016	92,582	92,487 (99.90)	663,157	3,074	4.64 (4.47,4.80)	65	3,009	4.54 (4.38, 4.70)
2017	97,405	97,341 (99.93)	646,794	2,947	4.56 (4.39,4.72)	57	2,890	4.47 (4.31, 4.63)
2018	96,641	96,629 (99.99)	625,310	2,872	4.59 (4.43,4.76)	69	2,803	4.48 (4.32, 4.65)
2019	94,734	94,717 (99.98)	610,140	2,591	4.25 (4.09, 4.41)	53	2,538	4.16 (4.00, 4.32)

Abbreviations: CI: confidence interval, GA: gestational age

Figure 1: Annual rates of brain injury occurring during or soon after birth in England (all gestational ages) 2012-2017 without exclusions; error bars indicate 95% confidence intervals



3.1.2 Rates of brain injuries in England: term infants

Numerator:	Annual number of term infants (≥ 37 weeks gestational age) who received at least one episode of care within a neonatal unit in England with a <i>brain injury occurring during or soon after birth</i> in England, with and without exclusions, obtained from the NNRD.
Denominator:	Annual number of term live births in England obtained from ONS Live Births by Gestational Age ¹¹ .

Table 2 presents the incidence and crude rates per 1,000 live term births of brain injuries in term infants, with and without exclusions of infants with conditions leading to brain injury prior to birth, for years 2012 to 2019. Figure 2 displays the crude rates of brain injuries in term infants per 1,000 live term births, prior to exclusions.


¹¹ Office for National Statistics, last accessed 14th December 2020, <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/12561livebirthsstillbirthsandneonataldeathsbygestationalageinengland2010to2018neonataldeathsand2010to2019livebirthsandstillbirths>

Table 2: Term infants in England (≥ 37 weeks gestational age) with a diagnosis of brain injury, before and after exclusion of infants with conditions leading to brain injury prior to birth.

Year	Term infants recorded in the NNRD	Term live births in England	Before exclusion of infants with conditions leading to brain injury prior to birth		After exclusion of infants with conditions leading to brain injury prior to birth		
			Term infants with brain injury	Rate of brain injuries per 1000 term live births (95%CI)	Exclusions	Term Infants with brain injury	Rate of brain injuries per 1000 term live births (95% CI)
2012	46,200	640,787	1714	2.67 (2.55, 2.80)	53	1661	2.59 (2.47, 2.72)
2013	47,935	612,816	1754	2.86 (2.73, 3.00)	41	1713	2.80 (2.67, 2.93)
2014	51,945	607,972	1848	3.04 (2.90, 3.18)	44	1804	2.97 (2.83, 3.11)
2015	55,045	609,076	1808	2.97 (2.83, 3.11)	37	1771	2.91 (2.78, 3.05)
2016	58,069	609,118	1760	2.89 (2.76, 3.03)	38	1722	2.83 (2.70, 2.96)
2017	62,558	593,339	1692	2.85 (2.72, 2.99)	42	1650	2.78 (2.65, 2.92)
2018	62,930	574,063	1,627	2.83 (2.70, 2.98)	40	1587	2.76 (2.63, 2.90)
2019	61,989	560,765	1,473	2.63 (2.50, 2.76)	34	1439	2.57 (2.44, 2.70)

Abbreviations: CI: confidence interval

Figure 2: Annual rates of brain injury occurring during or soon after birth in England in term infants (≥ 37 gestational weeks) 2012-2017 without exclusions; error bars indicate 95% confidence intervals



3.1.3 Rates of brain injuries in England: preterm infants

Numerator:	Annual number of preterm infants (<37 weeks gestational age) who received at least one episode of care within a neonatal unit in England with a <i>brain injury occurring during or soon after birth</i> in England, with and without exclusions, obtained from the NNRD.
Denominator:	Annual number of preterm live births in England obtained from ONS Live Births by Gestational Age ¹² .

Table 3 presents the incidence and crude rates per 1,000 live preterm births of brain injuries in preterm infants, with and without exclusions of infants with conditions leading to brain injury prior to birth, for years 2012 to 2019. Figure 3 displays the crude rates of brain injuries in preterm infants per 1,000 live preterm births, prior to exclusions.


¹² Office for National Statistics, last accessed 14th December 2020, <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/12561livebirthsstillbirthsandneonataldeathsbygestationalageinengland2010to2018neonataldeathsand2010to2019livebirthsandstillbirths>

Table 3: Preterm infants in England (<37 weeks gestational age) with a diagnosis of brain injury, before and after exclusion of infants with conditions leading to brain injury prior to birth.

Year*	Preterm infants recorded in the NNRD	Preterm live births in England	Before exclusion of infants with conditions leading to brain injury prior to birth		After exclusion of infants with conditions leading to brain injury prior to birth		
			Preterm infants with brain injury	Rate of brain injuries per 1000 term live births (95%CI)	Exclusions	Preterm infants with brain injury	Rate of brain injuries per 1000 preterm live births (95% CI)
2012	32,752	49,949	1236	24.75 (23.40, 26.16)	23	1213	24.28 (22.96, 25.69)
2013	32,264	48,844	1212	24.81 (23.46, 26.25)	21	1191	24.38 (23.04, 25.81)
2014	33,036	49,379	1249	25.29 (23.93, 26.74)	12	1237	25.05 (23.69, 26.49)
2015	33,740	50,308	1247	24.79 (23.45, 26.20)	17	1230	24.45 (23.12, 25.85)
2016	34,418	51,581	1314	25.47 (24.13, 26.89)	27	1287	24.95 (23.62, 26.35)
2017	34,783	51,307	1255	24.46 (23.14, 25.85)	15	1240	24.17 (22.86, 25.55)
2018	33,699	49,302	1245	25.25 (23.89, 26.69)	29	1216	24.66 (23.32, 26.09)
2019	33,028	47,488	1118	23.54 (22.20, 24.96)	19	1099	23.14 (21.81, 24.55)

Abbreviations: NNRD: National Neonatal Research Database; CI: confidence interval

Figure 3: Annual rates of brain injury occurring during or soon after birth in England in preterm infants (<37 gestational weeks) 2012-2017 without exclusions; error bars indicate 95% confidence intervals



3.1.4 Rates of brain injuries in England by medical condition

Numerators:	<p>Number of infants who received at least one instance of care within a neonatal unit in England with individual conditions included in the definition of <i>brain injuries occurring during or soon after birth</i> obtained from the NNRD:</p> <ol style="list-style-type: none"> 1. All gestational ages 2. Term births only (≥ 37 gestational weeks) 3. Preterm births only (< 37 gestational weeks)
Denominators:	<ol style="list-style-type: none"> 1. Annual number of live births in England obtained from ONS Birth Summary Tables¹³. 2. Annual number of term live births in England obtained from ONS Live Births by Gestational Age¹³ 3. Number of preterm live births in England obtained from ONS Live Births by Gestational Age¹³.

Tables 4 - 10 present the national rates of different medical conditions leading to brain injury at or soon after birth. Further explanation of way different medical conditions are defined, including the relevant NNRD data items, can be found in Appendix 1. It was possible for infants to be diagnosed with more than one condition leading to brain injury at or soon after birth, so the sum of total number of cases in tables 4 to 9 will not match totals in tables 1 to 3.

It is important to note that the the definition of hypoxic ischaemic encephalopathy (HIE) included within the definition of *brain injuries occurring during or soon after birth* is different from the definition used in other national reports (e.g. the National Neonatal Audit Programme audit measure *encephalopathy*¹⁴ and Each Baby Counts outcome *severe brain injury*¹⁵).

Low numbers of annual cases of *bilirubin encephalopathy* were reported. To ensure confidentiality and prevent disclosure of protected information cases of *bilirubin encephalopathy* are presented in aggregate over the six year period between 2012 and 2017.

¹³Office for National Statistics, last accessed 14th December 2020, <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/12561livebirthsstillbirthsandneonataldeathsbygestationalageinengland2010to2018neonataldeathsand2010to2019livebirthsandsstillbirths>

¹⁴ National Neonatal Audit Programme A guide to the 2018 audit measures; v1.2 published November 2018, last accessed 21st March 2019; <https://www.rcpch.ac.uk/sites/default/files/2018-11/NNAP%202018%20Audit%20Measures%20Guide%20v1.2%20FINAL.pdf>

¹⁵ Royal College of Obstetricians and Gynaecologists. Each Baby Counts: 2015 Full Report. London: RCOG, 2017, last accessed 21st March 2019; <https://www.rcog.org.uk/en/guidelines-research-services/audit-quality-improvement/each-baby-counts/ebc-2015-report/>



Table 4: Infants in England with hypoxic ischaemic encephalopathy (HIE)

HIE	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
Cases	1,128	1,161	1,257	1,243	1,204	1,171	1,164	983
Rate per 1000 live births (95% CI)	1.62 (1.53,1.72)	1.75 (1.65,1.85)	1.90 (1.80,2.01)	1.87 (1.77,1.98)	1.82 (1.72,1.92)	1.81 (1.71,1.92)	1.86 (1.76, 1.97)	1.61 (1.51,1.72)
Term cases	944	971	1,044	1,018	987	963	936	839
Preterm cases	184	190	213	225	217	208	228	144

Abbreviations: CI: confidence interval

Table 5: Infants in England with seizures

Seizures	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
Cases	1,445	1,432	1,360	1,249	1,223	1,159	1,127	1,016
Rate per 1000 live births (95% CI)	2.08 (1.98,2.19)	2.15 (2.05,2.27)	2.06 (1.95,2.17)	1.88 (1.78,1.99)	1.84 (1.74,1.95)	1.79 (1.69,1.90)	1.80 (1.70, 1.91)	1.67 (1.57, 1.77)
Term cases	1,065	1,036	1,009	919	866	845	806	725
Preterm cases	380	396	351	330	357	314	321	291

Abbreviations: CI: confidence interval



Table 6: Infants in England with intracranial haemorrhage

Intracranial haemorrhage	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
Cases	754	677	689	726	777	783	778	683
Rate per 1000 live births (95% CI)	1.09 (1.01,1.17)	1.02 (0.94,1.10)	1.04 (0.97,1.12)	1.09 (1.02,1.18)	1.17 (1.09,1.26)	1.21 (1.13,1.30)	1.24 (1.16, 1.33)	1.12 (1.04, 1.21)
Term cases	110	94	104	117	112	129	110	84
Rate per 10,000 term births (95% CI)	1.72 (1.42,2.07)	1.53 (1.25,1.88)	1.71 (1.41,2.07)	1.92 (1.60,2.30)	1.84 (1.53,2.21)	2.17 (1.83,2.58)	1.92 (1.59, 2.31)	1.50 (1.21, 1.86)
Preterm cases	644	583	585	609	665	654	668	599
Severe P/IVH <32 weeks GA	483	445	468	452	519	482	523	481
Rate of severe P/IVH per 1000 live births <32 weeks GA (95% CI)	60.35 (55.20,65.98)	57.72 (52.60,63.35)	61.13 (55.83,66.93)	58.27 (53.14,63.90)	61.76 (56.67,67.31)	58.64 (53.64,64.12)	67.29 (61.77, 73.31)	62.48 (57.13, 68.32)

Abbreviations: CI: confidence interval; P/IVH: periventricular/intraventricular haemorrhage; GA: gestational age



Table 7: Infants in England with central nervous system (CNS) infection

CNS infection	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
Cases	390	452	543	510	554	498	487	431
Rate per 1000 live births (95% CI)	0.56 (0.51,0.62)	0.68 (0.62,0.75)	0.82 (0.75,0.89)	0.77 (0.70,0.84)	0.84 (0.77,0.91)	0.77 (0.71,0.84)	0.78 (0.71, 0.85)	0.71 (0.64, 0.78)
Term cases	205	275	302	304	321	275	261	240
Preterm cases	185	177	241	206	233	223	226	191

Abbreviations: CI: confidence interval

Table 8: Infants in England with perinatal/neonatal stroke

Perinatal/ neonatal stroke	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
Cases	77	100	88	90	82	91	92	99
Rate per 1000 live births (95% CI)	0.11 (0.09,0.14)	0.15 (0.12,0.18)	0.13 (0.11,0.16)	0.14 (0.11,0.17)	0.12 (0.10,0.15)	0.14 (0.11,0.17)	0.15 (0.12, 0.18)	0.16 (0.13, 0.20)
Term cases	64	78	72	76	66	74	77	87
Preterm cases	13	22	16	14	16	17	15	12

Abbreviations: CI: confidence interval



Table 9: Infants in England with cystic periventricular leukomalacia (PVL)

Cystic PVL	Year							
	2012	2013	2014	2015	2016	2017	2018	2019
No. preterm cases	199	175	171	184	210	175	167	172
Rate per 1000 live births (95% CI)	0.3 (0.3, 0.3)	0.3 (0.2, 0.3)	0.3 (0.2, 0.3)	0.3 (0.2, 0.3)	0.3 (0.3, 0.4)	0.3 (0.2, 0.3)	0.3 (0.2, 0.3)	0.3 (0.2, 0.3)
No. cases at <34 weeks GA	186	175	157	176	201	171	161	165
Rate per 1000 live births <34 weeks GA (95% CI)	12.7 (11.0, 14.7)	12.5 (10.7, 14.4)	11.3 (9.7, 13.3)	12.4 (10.7, 14.4)	13.8 (12.0, 15.8)	11.8 (10.2, 13.7)	11.8 (10.1, 13.7)	12.3 (10.6, 14.4)

Abbreviations: CI: confidence interval; GA: gestational age



Table 10: Infants in England with bilirubin encephalopathy

Bilirubin Encephalopathy	Year
	2012 – 2019
All Cases	36
Annual rate per 100,000 live births (95% CI)	0.69 (0.50, 0.95)

Abbreviations: CI: confidence interval. Note: Due to low numbers of cases, data from 2012 to 2019 were merged.



Appendix 1: Definition of brain injuries occurring during or soon after birth using the National Neonatal Research Database (NNRD)

Appendix 1

Brain injury definition: National Neonatal Research Database (NNRD) data fields

Scope

- All babies admitted to a NHS neonatal unit in England
- Injury detected during neonatal unit stay to discharge

Data source

- Data extracted from the National Neonatal Research Database (NNRD) at the Neonatal Data Analysis Unit at Imperial College London.
- The NNRD contains a predefined set of variables (the Neonatal Data Set, an authorised NHS Information Standard) extracted at regular intervals from the Electronic Patient Record of every admission to a NHS neonatal unit in England, Wales, and Scotland, cleaned and merged across multiple patient episodes, to create a single data file for each patient.



Appendix 1: Definition of brain injuries occurring during or soon after birth using the National Neonatal Research Database (NNRD)

Condition	Data items
HIE	<p>Any of the following recorded in any <i>Diagnosis</i> field:</p> <ol style="list-style-type: none"> 1. Severe Hypoxic Ischaemic Encephalopathy (HIE) 2. Severe Neonatal Encephalopathy 3. Grade 3 Hypoxic Ischaemic Encephalopathy (HIE) 4. Moderate Hypoxic Ischaemic Encephalopathy (HIE) 5. Moderate Neonatal Encephalopathy 6. Grade 2 Hypoxic Ischaemic Encephalopathy (HIE) <p style="text-align: center;">OR</p> <p>The following recorded in <i>daily care neurology</i> field:</p> <ol style="list-style-type: none"> 1. Therapeutic hypothermia induced - for 2 or more consecutive days
Intracranial haemorrhage	<p>Any of the following recorded in any <i>Diagnosis</i> field:</p> <ol style="list-style-type: none"> 1. Subdural haemorrhage due to birth injury 2. Cerebral haemorrhage due to birth injury 3. Traumatic intraventricular haemorrhage 4. Subarachnoid haemorrhage due to birth injury 5. Subarachnoid haemorrhage 6. Tentorial tear due to birth injury 7. Intracranial laceration and haemorrhage due to birth injury 8. Large intraventricular haemorrhage (IVH Grade 3) 9. Intraventricular haemorrhage/parenchymal 10. Parenchymal haemorrhage 11. haemorrhage (IVH Grade 4) 12. Intracranial Haemorrhage (unknown or unspecified cause) 13. Intracerebral haemorrhage 14. Intracerebral haemorrhage (term infant) 15. Intraventricular haemorrhage (perinatal) 16. Post-haemorrhagic hydrocephalus <p style="text-align: center;">OR</p> <p>Any of the following recorded in any <i>cranial ultrasound findings</i> field:</p> <ol style="list-style-type: none"> 1. Large intraventricular haemorrhage (IVH Grade 3) 2. Intraventricular haemorrhage/parenchymal haemorrhage (IVH Grade 4) 3. Parenchymal haemorrhage <p style="text-align: center;">OR</p> <p>Any of the following recorded in any <i>procedure field</i>:</p> <ol style="list-style-type: none"> 1. Ventriculoperitoneal or other ventricular shunt 2. External ventricular drain 3. Ventricular drain with reservoir 4. Insertion of ventricular peritoneal shunt.



Appendix 1: Definition of brain injuries occurring during or soon after birth using the National Neonatal Research Database (NNRD)

	<ol style="list-style-type: none"> 5. Insertion of Rickham reservoir 6. Insertion of ventriculo-atrial CSF shunt 7. Insertion of ventriculo-peritoneal CSF shunt 8. Creation of ventriculoperitoneal shunt
Preterm white matter injury	<p>Any of the following recorded in any <i>Diagnosis</i> field:</p> <ol style="list-style-type: none"> 1. Cystic periventricular leukomalacia <p style="text-align: center;">OR</p> <p>Any of the following recorded in any <i>cranial ultrasound findings</i> field:</p> <ol style="list-style-type: none"> 1. Cystic periventricular leucomalacia
Perinatal stroke	<p>Any of the following recorded in any <i>Diagnosis at discharge</i> field:</p> <ol style="list-style-type: none"> 1. Neonatal stroke 2. Infarction: Middle cerebral artery (stroke) 3. Cerebrovascular accident (stroke) 4. Cerebral venous thrombosis 5. Neonatal cerebral ischaemia
Central nervous system infection	<p>Any of the following diagnosis codes recorded in the <i>Diagnosis</i> field:</p> <ol style="list-style-type: none"> 1. Bacterial meningitis 2. Viral meningitis 3. Meningitis – streptococcal 4. Meningitis – bacterial (specific organism) 5. Meningitis – bacterial (unknown or unspecified organism) 6. Meningitis – Candida 7. Candida encephalitis 8. Congenital herpes infection <p>Any pathogen recorded in the <i>suspected infection data</i> field <i>Pathogen in CSF</i></p>
Kernicterus	<p>Any of the following diagnoses recorded in any <i>Diagnosis</i> field:</p> <ol style="list-style-type: none"> 1. Bilirubin encephalopathy (immune) 2. Kernicterus (unspecified or unknown cause) 3. Kernicterus
Seizures	<p>Any of the following recorded in any <i>daily care neurology</i> field:</p> <ol style="list-style-type: none"> 1. Seizure occurred



Appendix 1: Definition of brain injuries occurring during or soon after birth using the National Neonatal Research Database (NNRD)

Exclusions

Data will be presented before and after exclusion of infants with **seizures** (condition defined as above) AND the following diagnosis codes recorded in the *Diagnosis* field during their their neonatal unit admission.

Condition

Congenital encephalopathies

Data items

Episodic variables: Any of the following diagnosis codes entered into any *Diagnosis* field:

1. Congenital neuropathy (unknown or unspecified cause)
2. Congenital myopathy
3. Mitochondrial myopathy
4. Congenital Central Hypoventilation Syndrome (CCHS)
5. Congenital hypertonia
6. Congenital hypotonia - floppy
7. Benign familial neonatal seizures
8. Inborn error of metabolism (description required)
9. Myotonic dystrophy requiring endotracheal intubation and assisted ventilation
10. Disorder of branch chain amino acid metabolism
11. Disorders of fatty acid metabolism
12. Disorders of fatty acid metabolism: carnitine metabolism
13. Disorder of glycine metabolism
14. Disorder of glycine metabolism: Non ketotic hyperglycinaemia
15. Hyperammonaemia of the newborn
16. Disorders of lysine and hydroxylysine metabolism
17. Disorders of ornithine metabolism
18. Disorders of pyruvate metabolism and gluconeogenesis
19. Disorder of carbohydrate metabolism (unknown or unspecified cause)
20. Down Syndrome (Trisomy 21)
21. Trisomy 21
22. Edwards Syndrome (Trisomy 18)
23. Trisomy 18
24. Patau Syndrome (Trisomy 13)
25. Trisomy 13

Congenital infections

Episodic variables: Any of the following diagnosis codes entered into any *Diagnosis* field:

1. Congenital viral disease (specify)
2. Syphilis - latent congenital
3. Congenital rubella syndrome



Appendix 1: Definition of brain injuries occurring during or soon after birth using the National Neonatal Research Database (NNRD)

Congenital brain
abnormalities

4. Congenital cytomegalovirus infection
5. Congenital herpes [herpes simplex] infection
6. Other congenital viral diseases
7. Congenital viral disease (unknown or unspecified cause)
8. Congenital toxoplasmosis

Episodic variables: Any of the following diagnosis codes entered into any *Diagnosis* field:

1. Malformations of aqueduct of Sylvius
2. Atresia of foramina of Magendie and Luschka
3. Atresia of foramina of Magendie and Luschka
4. Other congenital hydrocephalus
5. X linked congenital hydrocephalus
6. Congenital hydrocephalus (unknown or unspecified cause)
7. Congenital hydrocephalus
8. Lissencephaly
9. Vermal agenesis
10. Septum pelucidum absence – congenital
11. Congenital malformations of corpus callosum
12. Arhinencephaly
13. Holoprosencephaly
14. Other reduction deformities of brain
15. Septo-optic dysplasia
16. Other specified congenital malformations of brain
17. Congenital malformation of brain (unknown or unspecified cause)
18. Other congenital malformations of brain
19. Cervical spina bifida with hydrocephalus
20. Thoracic spina bifida with hydrocephalus
21. Spina bifida (unknown or unspecified cause)
22. Spina bifida
23. Amyelia
24. Hypoplasia and dysplasia of spinal cord
25. Other specified congenital malformations of spinal cord
26. Congenital malformation of spinal cord (unknown or unspecified cause)
27. Nerve palsies – congenital
28. Arnold-Chiari syndrome
29. Other specified congenital malformations of nervous system
30. Congenital malformation of nervous system (unknown or unspecified cause)
31. Other congenital malformations of nervous system
32. Congenital Hydrocephalus



